

THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

November 10, 2004

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE.

APPLICATION NUMBER: 60/560,719

FILING DATE: *October 07, 2003*

RELATED PCT APPLICATION NUMBER: *PCT/US04/32986*

Certified by

Jon W Dudas



Acting Under Secretary of Commerce
for Intellectual Property
and Acting Director of the U.S.
Patent and Trademark Office

HAMILTON, BROOK, SMITH & REYNOLDS, P.C.

UTILITY PATENT APPLICATION TRANSMITTAL <small>(Only for new nonprovisional applications under 37 C.F.R. 1.53(b))</small>	Attorney Docket No.	3268.1007-000
	First Named Inventor or Application Identifier	Yousef Al-Abed
	Express Mail Label No.	EV 214947007 US

Title of Invention	ANTI-INFLAMMATORY COMPOUNDS
-----------------------	-----------------------------

APPLICATION ELEMENTS See MPEP chapter 600 concerning utility patent application contents.	ADDRESS TO: Mail Stop Patent Application Commissioner for Patents P.O. Box 1450 Alexandria, VA 22313-1450
---	--

1. <input type="checkbox"/> Fee Transmittal Form <i>(Submit an original, and a duplicate for fee processing)</i> 2. <input checked="" type="checkbox"/> Specification Total Pages [50] <i>(preferred arrangement set forth below)</i> - Descriptive title of the invention - Cross References to Related Applications - Statement Regarding Fed sponsored R & D - Reference to sequence listing, a table, or a computer program listing appendix - Background of the Invention - Brief Summary of the Invention - Brief Description of the Drawings <i>(if filed)</i> - Detailed Description - Claim(s) - Abstract of the Disclosure 3. <input checked="" type="checkbox"/> Drawing(s) (35 U.S.C. 113) Total Sheets [6] <input type="checkbox"/> Fig. of the Drawings for Publication <input type="checkbox"/> <input checked="" type="checkbox"/> No Figure to be Published 4. <input type="checkbox"/> Oath or Declaration Total Pages [] a. <input type="checkbox"/> Newly executed (original or copy) b. <input type="checkbox"/> Copy from a prior application (37 C.F.R. 1.63(d)) <i>(for continuation/divisional with Box 17 completed)</i> i. <input type="checkbox"/> <u>DELETION OF INVENTOR(S)</u> Signed statement attached deleting inventor(s) named in the prior application, see 37 C.F.R. 1.63(d)(2) and 1.33(b). 5. <input type="checkbox"/> CD-ROM or CD-R in duplicate, large table or Computer Program <i>(Appendix)</i>	6. <input type="checkbox"/> Nucleotide and/or Amino Acid Sequence Submission (if applicable, all necessary) a. <input type="checkbox"/> Computer Readable Form b. <input type="checkbox"/> Paper Copy (identical to computer copy) [] Pages c. <input type="checkbox"/> Statements verifying identity of above copies <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="text-align: center;">ACCOMPANYING APPLICATION PARTS</td> </tr> <tr> <td> 7. <input type="checkbox"/> Assignment Papers (cover sheet & documents) <input checked="" type="checkbox"/> Assignee - North Shore-Long Island Jewish Research Institute Manhasset, NY 8. <input type="checkbox"/> Power of Attorney <input type="checkbox"/> 37 C.F.R. 3.73(b) Statement 9. <input type="checkbox"/> English Translation Document <i>(if applicable)</i> 10. <input type="checkbox"/> Information Disclosure <input type="checkbox"/> Copies of IDS Citations Statement (IDS)/PTO-1449 11. <input type="checkbox"/> Preliminary Amendment 12. <input checked="" type="checkbox"/> Return Receipt Postcard 13. <input type="checkbox"/> Small Entity Statement(s) 14a. <input type="checkbox"/> Foreign Priority Claim under 35 U.S.C. § 119 or 365 14b. <input type="checkbox"/> Certified Copy of Priority Document(s) 15. <input type="checkbox"/> Nonpublication Request <i>(check parent application)</i> 16. <input type="checkbox"/> Other _____ </td> </tr> </table>	ACCOMPANYING APPLICATION PARTS	7. <input type="checkbox"/> Assignment Papers (cover sheet & documents) <input checked="" type="checkbox"/> Assignee - North Shore-Long Island Jewish Research Institute Manhasset, NY 8. <input type="checkbox"/> Power of Attorney <input type="checkbox"/> 37 C.F.R. 3.73(b) Statement 9. <input type="checkbox"/> English Translation Document <i>(if applicable)</i> 10. <input type="checkbox"/> Information Disclosure <input type="checkbox"/> Copies of IDS Citations Statement (IDS)/PTO-1449 11. <input type="checkbox"/> Preliminary Amendment 12. <input checked="" type="checkbox"/> Return Receipt Postcard 13. <input type="checkbox"/> Small Entity Statement(s) 14a. <input type="checkbox"/> Foreign Priority Claim under 35 U.S.C. § 119 or 365 14b. <input type="checkbox"/> Certified Copy of Priority Document(s) 15. <input type="checkbox"/> Nonpublication Request <i>(check parent application)</i> 16. <input type="checkbox"/> Other _____
ACCOMPANYING APPLICATION PARTS			
7. <input type="checkbox"/> Assignment Papers (cover sheet & documents) <input checked="" type="checkbox"/> Assignee - North Shore-Long Island Jewish Research Institute Manhasset, NY 8. <input type="checkbox"/> Power of Attorney <input type="checkbox"/> 37 C.F.R. 3.73(b) Statement 9. <input type="checkbox"/> English Translation Document <i>(if applicable)</i> 10. <input type="checkbox"/> Information Disclosure <input type="checkbox"/> Copies of IDS Citations Statement (IDS)/PTO-1449 11. <input type="checkbox"/> Preliminary Amendment 12. <input checked="" type="checkbox"/> Return Receipt Postcard 13. <input type="checkbox"/> Small Entity Statement(s) 14a. <input type="checkbox"/> Foreign Priority Claim under 35 U.S.C. § 119 or 365 14b. <input type="checkbox"/> Certified Copy of Priority Document(s) 15. <input type="checkbox"/> Nonpublication Request <i>(check parent application)</i> 16. <input type="checkbox"/> Other _____			

17. If a CONTINUING APPLICATION, check appropriate box; supply the requisite information. <input type="checkbox"/> Continuation <input type="checkbox"/> Divisional <input type="checkbox"/> Continuation-in-part (CIP) of prior application No.: Prior application information: Examiner: Group Art Unit:			
The entire disclosure of the prior application is considered a part of the disclosure of the accompanying application and is hereby incorporated by reference. <i>(Add standard Related Applications section with incorporation by reference to specification or update same)</i>			

18. CORRESPONDENCE ADDRESS					
NAME	Customer No. 021005				
	HAMILTON, BROOK, SMITH & REYNOLDS, P.C.				
ADDRESS	530 Virginia Road, P.O. Box 9133				
CITY	Concord	STATE	MA	ZIP CODE	01742-9133
COUNTRY	USA	TELEPHONE	(978) 341-0036	FAX	(978) 341-0136

Signature	<i>Alexander Akhiezer</i>	Date	10/2/03
Submitted by Typed or Printed Name	Alexander Akhiezer	Reg. Number	54,617

ODMA\HODMA\Manage;425472;1

Copied from 10680715 on 07/26/2004

- 1 -

Date: <u>10/7/03</u>	Express Mail Label No. <u>EV 2149470711</u>
----------------------	---

Inventors: Yousef Al-Abed and Kevin J. Tracey

Attorney's Docket No.: 3268.1007-000

ANTI-INFLAMMATORY COMPOUNDS

GOVERNMENT SUPPORT

The invention was supported, in whole or in part, by a grant N00178-01-C3058 from DARPA. The Government has certain rights in the invention.

5 BACKGROUND OF THE INVENTION

Inflammation is often induced by proinflammatory cytokines, such as tumor necrosis factor (TNF), interleukin (IL)-1 α , IL-1 β , IL-6, platelet-activating factor (PAF), macrophage migration inhibitory factor (MIF), HMGB1 and other compounds. These proinflammatory cytokines are produced by several different cell types, most importantly immune cells (for
10 example, monocytes, macrophages and neutrophils), contributing to various disorders during the early stages of an inflammatory cytokine cascade.

The early pro-inflammatory cytokines (e.g., TNF, IL-1, etc.) mediate inflammation, and induce the late release of high mobility group-1 (HMG1) (also known as HMG-1 and HMGB1), a protein that accumulates in serum and mediates delayed lethality and further
15 induction of early pro-inflammatory cytokines.

Tumor Necrosis Factor (TNF) isoforms α and β are soluble proteins produced by monocytes and macrophages in response to endotoxin or other stimuli. Cells other than monocytes or macrophages also make TNF α . For example, human non-monocytic tumor cell lines produce TNF α . (For recent reviews on TNF family, see Lin A, *et al.*, Aging Cell.
20 2002 Dec;1(2):112-6 and Baugh J. *et al.*, Curr Opin Drug Discov Devel. 2001 Sep;4(5):635-50.)

Upon binding to its cognate receptor, TNF α activates both apoptosis and NF- κ B-dependent survival pathways. TNF causes pro-inflammatory actions which result in tissue injury, such as inducing pro-coagulant activity on vascular endothelial cells, increasing the adherence of neutrophils and lymphocytes, and stimulating the release of platelet activating factor from macrophages, neutrophils and vascular endothelial cells. (Pober, *et al.*, *J. Immunol.* 136:1680 (1986); Pober, *et al.*, *J. Immunol.* 138:3319 (1987); Camussi, *et al.*, *J. Exp. Med.* 166:1390 (1987)).

TNF is also associated with infections, immune disorders, neoplastic, autoimmune and graft-versus host pathologies. TNF can mediate cachexia in cancer, infectious pathology, and other catabolic states. (Kern, *et al.*, *J. Parent. Enter. Nutr.* 12:286-298 (1988)). TNF also plays a central role in gram-negative sepsis and endotoxic shock. Endotoxin strongly activates monocyte/macrophage production and secretion of TNF and other cytokines. (Michie, *et al.*, *Br. J. Surg.* 76:670-671 (1989); Debets, *et al.*, *Second Vienna Shock Forum*, p.463-466 (1989); Simpson, *et al.*, *Crit. Care Clin.* 5:27-47 (1989); Kornbluth, *et al.*, *J. Immunol.* 137:2585-2591 (1986)).


HMGB1 is a member of a family of DNA-binding proteins termed high mobility group (HMG) and has been implicated as a cytokine mediator of delayed lethality in endotoxemia (Wang *et al.*, *Science* 285: 248-251, 1999; WO 00/47104). That work demonstrated that bacterial endotoxin (lipopolysaccharide (LPS)) activates monocytes/macrophages to release HMGB1 as a late response to activation, resulting in elevated serum HMGB1 levels that are toxic. Antibodies against HMGB1 prevent lethality of endotoxin even when antibody administration is delayed until after the early cytokine response. (*ibid.*) HMGB1 has also been implicated in mediating other inflammatory disorders such as endotoxin-induced lung edema, sepsis and hemorrhagic shock. (See, *e.g.*, U.S. Patents Nos. 6,448,223 and 6,468,533.)

There is a need for new therapeutic agents which suppress the production, release and/or activity of inflammatory cytokines such as TNF and HMGB1 and which therefore can be used in treatment of inflammatory disorders.

SUMMARY OF THE INVENTION

It has now been discovered that certain derivatives of isoxazole, isoxazolidine, isothiazole and isothiazolidine compounds inhibit production and release of certain proinflammatory cytokines such as TNF and HMGB1 (see Examples 1, 2 and 5). Based on this discovery, methods of treating a subject with an inflammatory cytokine-mediated disorder as well as pharmaceutical compositions and compositions of matter that include the anti-inflammatory compounds of the present invention are disclosed.

In one embodiment, the present invention is a compound of Formula (I) or a pharmaceutically acceptable salt thereof:




Ar₁ and Ar₂ are independently a monocyclic six-member optionally substituted heteroaryl group.

A₁ is =N- or -NR^a- and A₂ is O or S; R^a is H or C1-C6 alkyl.

R₁ is selected from -H, C1-C6 alkyl, phenyl, C1-C6 haloalkyl, halogen, -OH, -OR^b, C1-C6 hydroxyalkyl, C1-C6 alkoxyalkyl, -O(C1-C6 haloalkyl), -SH, -SR^b, -NO₂, -CN, -NR^bCO₂ R^b, -NR^bC(O) R^b, -CO₂R^b, -C(O)R^b, -C(O)N(R^b)₂, -OC(O)R^b and -NR^bR^b.

Each R^b is H or a C1-C6 alkyl group.


In another embodiment, the present invention is a pharmaceutical composition comprising a pharmaceutically acceptable carrier or diluent and a compound represented by Formula (I a) or a pharmaceutically acceptable salt thereof:



Ar₁, Ar₂, A₁, A₂, R₁ and its substituents are defined above for Formula (I).

In another embodiment, the present invention is a pharmaceutical composition comprising a pharmaceutically acceptable carrier or diluent and a compound represented by Formula (VII) or a pharmaceutically acceptable salt thereof:

5



Ar is an optionally substituted, monocyclic, six-member heteroaryl.

A₁ is =N- or -NR^a- and A₂ is O or S.


10 R₁ is -H, C1-C6 alkyl, phenyl, C1-C6 haloalkyl, halogen, -OH, -OR^b, C1-C6 hydroxyalkyl, C1-C6 alkoxyalkyl, -O(C1-C6 haloalkyl), -SH, -SR^b, -NO₂, -CN, -NR^bCO₂ R^b, -NR^bC(O) R^b, -CO₂R^b, -C(O)R^b, -C(O)N(R^b)₂, -OC(O)R^b or -NR^bR^b.

Each R^a is -H or C1-C6 alkyl and each R^b is -H or a C1-C6 alkyl group.

Ring D is optionally substituted with zero, one or more substituents other than amide and is not an alkylphenol.

15 In another embodiment, the present invention is a method of treating a subject with an inflammatory cytokine mediated disorder comprising administering to the subject a therapeutically effective amount of a compound represented by Formula (I a) or a pharmaceutically acceptable salt thereof.

20 In another embodiment, the present invention is a method of treating a subject with an inflammatory cytokine mediated disorder comprising administering to the subject a therapeutically effective amount of a compound represented by Formula (VII a) or a pharmaceutically acceptable salt thereof:



Ar, A₁, A₂, R₁ and ring **D** and its substituents are as defined above for Formula (VII).

The present invention offers the advantage of providing individuals in need of treatment for inflammatory conditions with effective pharmaceutical agents.

5 BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is plot of the concentration of TNF, released by LPS-stimulated RAW cells, as a function of concentration of the compound of Formula (VI f) in the RAW cell culture. A plot of concentration of TNF as a function of nicotine concentration is included as a control.

FIG. 2 is a side-by-side comparison of plots representing HMGB-1 release by LPS-stimulated RAW cells, as a function of concentration of either the compound of Formula (VI f) or nicotine in the RAW cell culture. The release is measured as percentage of LPS-stimulated HMGB-1 released by untreated cells.

FIG. 3 is a table listing IC₅₀ values for the compounds of Formulae (VI f), (VI b), (XII b), (VI e), (VI g) and (XII a).

FIG. 4 is a table showing the number of mice, by days, protected by the compound of Formula (VI f) against septic shock induced according to CLP model.

FIG. 5 is a side-by-side comparison of plots representing TNF production by LPS-stimulated human macrophages treated by either the compound of Formula (VI f), (VI d) or nicotine.




FIG. 6 is a plot of TNF α production by LPS-stimulated RAW cells in the presence of either the compound of Formula (VI f) alone or in combination with α -bingarotoxin.

DETAILED DESCRIPTION OF THE INVENTION

The present invention encompasses pharmaceutical agents, pharmaceutical compositions and methods of treatment of cytokine-mediated inflammatory disorders, particularly, the disorders mediated by pro-inflammatory cytokines such as tumor necrosis factor (TNF), high mobility group protein HMGB1, interleukin (IL)-1 α , IL-1 β , IL-6, platelet-activating factor (PAF) and macrophage migration inhibitory factor (MIF). The compounds and methods of the invention are particularly useful for treatment of inflammatory disorders mediated by TNF- α and HMGB1.

In one embodiment, the present invention is a compound represented by Formula (I) or a pharmaceutically acceptable salt thereof. The variables of Formula (I) are as described above.

In a preferred embodiment, the compound of Formula (I) is represented by Formulae (II), (III) or (IV):








The variables for Formulae (II) to (IV) are described below.


B₁ through B₅ and D₁ through D₅ are independently N or CR^c, provided that from one to three of B₁ through B₅ and from one to three of D₁ through D₅ are N. Each R^c is independently any suitable substituent as described below for a heteroaryl group. R₁ is as described above for Formula (I).

More preferably, in structural Formulae (II) to (IV), R_1 is -H or a C1-C3 alkyl, optionally substituted with a halogen or a hydroxyl, and/or R^c is -H, halogen, $-\text{NO}_2$, $-\text{CN}$, C1-C3 alkyl, C1-C3 haloalkyl, C1-C3 hydroxyalkyl, C1-C3 alkoxyalkyl, $-\text{N}(\text{R}^d)_2$, $-\text{NR}^d\text{C}(\text{O})\text{R}^d$, or $-\text{C}(\text{O})\text{N}(\text{R}^d)_2$. R^d is H or C1-C3 alkyl.

- 5 In a more preferred embodiment, the anti-inflammatory compounds of the present invention are represented by Formulae (V a) through (V i):




- 15 R' and R'' are independently -H, halogen, $-\text{NO}_2$, $-\text{CN}$, C1-C3 alkyl, C1-C3 haloalkyl, C1-C3 hydroxyalkyl, C1-C3 alkoxyalkyl, $-\text{N}(\text{R}^d)_2$, $-\text{NR}^d\text{C}(\text{O})\text{R}^d$, or $-\text{C}(\text{O})\text{N}(\text{R}^d)_2$. R^d is as defined above. Specific examples of the compounds of the present invention are:



In a particularly preferred embodiment, the compounds are represented by Formula (VI d) or (VI f).

In another embodiment, the present invention is a pharmaceutical composition comprising a pharmaceutically acceptable carrier or diluent and any of the compounds represented by Formulae (I), (I a), (II), (III), (IV), (V a) through (V i) and (VI a) through (V i) or pharmaceutically acceptable salts thereof.



In one embodiment the disclosed invention is a pharmaceutical composition comprising a compound of Formula (I a). Preferably, the compound is represented by Formula (II a):



where variables B₁ through B₅, D₁ through D₅, A₁, A₂ and R₁ are defined above for Formula (I).

In another embodiment, the present invention is a pharmaceutical composition comprising a pharmaceutically acceptable carrier or diluent and a compound represented by Formula (VII). The variables of Formula (VII) are as described above.

In preferred embodiments, the compounds of Formula (VII) are represented by Formulae (VIII) and (IX):



The variables of Formulae (VIII) and (IX) are as follows.


A₁, A₂ and R₁ are as defined above for Formula (VII).

B₁ through B₅ are independently N or CR^c, provided that from one to three of B₁ through B₅ are N.

Each R^c is independently any suitable substituent as described below for a heteroaryl group.

Ring D is optionally substituted with zero, one or more substituent R₂. Each R₂ is independently any suitable substituent described below for an aryl group.

More preferably, the compounds of Formula (VII) are represented by Formulae (X) and (XI a) to (XI c):




5

where one of B_1 through B_3 is N, n is 0, 1 or 2 and m is 0, 1 or 2.

Preferably, in Formulae (VIII) to (XI a) through (XI c) R^c is -H, halogen, $-\text{NO}_2$, $-\text{CN}$, C1-C3 alkyl, C1-C3 haloalkyl, C1-C3 hydroxyalkyl, C1-C3 alkoxyalkyl, $-\text{N}(\text{R}^d)_2$, $-\text{NR}^d\text{C}(\text{O})\text{R}^d$, or $-\text{C}(\text{O})\text{N}(\text{R}^d)_2$ and R_2 is -H, halogen, $-\text{NO}_2$, $-\text{CN}$, C1-C3 alkyl, C1-C3 haloalkyl, C1-C3 hydroxyalkyl or C1-C3 alkoxyalkyl. R^d is H or C1-C3 alkyl.


10

Specific examples of the compounds suitable for using in pharmaceutical compositions of the present invention are



15

In one embodiment, the compounds of Formula (VII a), suitable for using in the methods of the present invention, are represented by Formula (VIII a):



where B₁ through B₅ and the substituents thereof are defined above for Formula (VIII).

In another embodiment, the compounds suitable for use in the methods of the present invention are represented by Formulae (I), (I a), (II), (II a) (III), (IV), (V a) to (V i), (VI a) to (VI i), (VII), (VII a), (VIII), (IX), (X), (XI a) through (XI c), (XII a) and (XII b) as defined above.

The term "heteroaryl", as used herein, refers to aromatic groups containing one or more heteroatoms (O, S, or N). A heteroaryl group of the present invention is a monocyclic six-member group. The heteroaryl groups of this invention can also include ring systems substituted with one or more oxo moieties. Examples of heteroaryl groups include, but are not limited to, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, pyridazinyl, pyridazinyl and triazinyl.

Suitable substituents on a heteroaryl group (including heteroaryl groups represented by Ar, Ar₁ and Ar₂) are those that do not substantially interfere with the pharmaceutical activity of the disclosed compound. A heteroaryl may have one or more substituents, which can be identical or different. Examples of suitable substituents for a substitutable carbon atom in a heteroaryl group include -H, C1-C6 alkyl, halogen, C1-C6 haloalkyl, -R^o, -OH, -OR^o, -O(C1-C6 haloalkyl), -SH, -SR^o, -NO₂, -CN, -NH₂, -NHCO₂R^o, -NHC(O)H, -NHC(O)R^o, -NR^oC(O)R^o, -CO₂H, -CO₂R^o, -C(O)H, -C(O)R^o, -C(O)NHR^o, -C(O)N(R^o)₂, -OC(O)R^o, -S(O)₂R^o, -SO₂NH₂, -S(O)R^o, -NHSO₂R^o, or a C1-C6 alkyl group substituted with R^o, -OH, -OR^o, -SH, -SR^o, -NO₂, -CN, -NHCO₂R^o, -NHC(O)H, -NHC(O)R^o, -CO₂H, -CO₂R^o, -C(O)H, -C(O)R^o, -C(O)NHR^o, -OC(O)R^o, -S(O)₂R^o, -SO₂NH₂, -S(O)R^o or -NHSO₂R^o. R^o is independently, C1-C6 alkyl, aryl or heteroaryl group and wherein the aryl or heteroaryl group represented by R^o is optionally substituted with one or more halogen, methyl or methoxy groups.

The term "aryl", as used herein, refers to a carbocyclic aromatic group. Examples of aryl groups include, but are not limited to phenyl and naphthyl.

A substituted aryl group can have one or more substituents which can be the same or different. Suitable substituents for a substituted aryl group, including ring **D**, typically represented herein as "R₂", are as defined above for a heteroaryls, provided that the substituents on ring **D** are other than amide and that ring **D** is not an alkylphenol. As used herein, the term "amide" refers to a -C(O)NHR^o group, where R^o is defined above for heteroaryl groups. As used herein, the term "alkylphenol" refers to a six-member monocyclic aryl substituted with one hydroxyl groups and one or more alkyls. Examples of suitable substituents for an aryl include -H, halogen, C1-C6 haloalkyl, -R^o, -OH, -OR^o, -O(C1-C6 haloalkyl), -SH, -SR^o, -NO₂, -CN, -NHCO₂R^o, -NHC(O)H, -NHC(O)R^o, -CO₂H, -CO₂R^o, -C(O)H, -C(O)R^o, -OC(O)R^o, -S(O)₂R^o, -SO₂NH₂, -S(O)R^o, -NHSO₂R^o, or a C1-C6 alkyl group substituted with R^o, -OH, -OR^o, -SH, -SR^o, -NO₂, -CN, -NHCO₂R^o, -NHC(O)H, -NHC(O)R^o, -CO₂H, -CO₂R^o, -C(O)H, -C(O)R^o, -C(O)NHR^o, -OC(O)R^o, -S(O)₂R^o, -SO₂NH₂, -S(O)R^o or -NHSO₂R^o. R^o is as defined above for heteroaryl groups.

The term "alkyl", as used herein, unless otherwise indicated, includes straight, branched or cyclic saturated monovalent hydrocarbon radicals, typically C1-C10, preferably C1-C6. Examples of alkyl groups include, but are not limited to, methyl, ethyl, propyl, isopropyl, and t-butyl.

The term "haloalkyl", as used herein, includes an alkyl substituted with one or more F, Cl, Br, or I, wherein the alkyl is defined above.

The terms "alkoxy", as used herein, means an "alkyl-O-" group, wherein alkyl, is defined above.

Pharmaceutically acceptable salts of the anti-inflammatory compounds disclosed herein can be used to practice the present invention. As used herein, a "pharmaceutically acceptable salt" of the disclosed compound is an ionic bond-containing product of reacting a compound of the invention with either an acid or a base, suitable for administering to a subject. For example, an acid salt of a compound containing an amine or other basic group can be obtained by reacting the compound with a suitable organic or inorganic acid, such as hydrogen chloride, hydrogen bromide, acetic acid, perchloric acid and the like. Other

examples of such salts include hydrochlorides, hydrobromides, sulfates, methanesulfonates, nitrates, maleates, acetates, citrates, fumarates, tartrates (e.g. (+)-tartrates, (-)-tartrates or mixtures thereof including racemic mixtures), succinates, benzoates and salts with amino acids such as glutamic acid. Salts can also be formed with suitable organic bases when the compound comprises an acid functional group such as $-\text{COOH}$ or $-\text{SO}_3\text{H}$. Such bases suitable for the formation of a pharmaceutically acceptable base addition salts with compounds of the present invention include organic bases that are nontoxic and strong enough to react with the acid functional group. Such organic bases are well known in the art and include amino acids such as arginine and lysine, mono-, di-, and triethanolamine, choline, mono-, di-, and trialkylamine, such as methylamine, dimethylamine, and trimethylamine, guanidine, N-benzylphenethylamine, N-methylglucosamine, N-methylpiperazine, morpholine, ethylenediamine, tris(hydroxymethyl)aminomethane and the like.

As used herein, a "pharmaceutical composition" is a formulation comprising the disclosed compounds and a pharmaceutically acceptable diluent or carrier, in a form suitable for administration to a subject. The pharmaceutical composition can be in bulk or in unit dosage form. The unit dosage form can be in any of a variety of forms, including, for example, a capsule, an IV bag, a tablet, a single pump on an aerosol inhaler, or a vial. The quantity of active ingredient (i.e., a formulation of the disclosed compound or salts thereof) in a unit dose of composition is an effective amount and may be varied according to the particular treatment involved. It may be appreciated that it may be necessary to make routine variations to the dosage depending on the age and condition of the patient. The dosage will also depend on the route of administration. A variety of routes are contemplated, including topical, oral, pulmonary, rectal, vaginal, parenteral, including transdermal, subcutaneous, intravenous, intramuscular, intraperitoneal and intranasal.

The compounds described herein, and the pharmaceutically acceptable salts thereof can be used in pharmaceutical preparations in combination with a pharmaceutically acceptable carrier or diluent. Suitable pharmaceutically acceptable carriers include inert solid fillers or diluents and sterile aqueous or organic solutions. The compounds will be present in such pharmaceutical compositions in amounts sufficient to provide the desired

dosage amount in the range described herein. Techniques for formulation and administration of the compounds of the instant invention can be found in *Remington: the Science and Practice of Pharmacy*, 19th edition, Mack Publishing Co., Easton, PA (1995).

As used herein, a "subject" includes mammals, e.g., humans, companion animals (e.g., dogs, cats, birds and the like), farm animals (e.g., cows, sheep, pigs, horses, fowl and the like) and laboratory animals (e.g., rats, mice, guinea pigs and the like). In a preferred embodiment of the disclosed methods, the subject is human.

As used herein, a "therapeutically effective amount" of a compound of the disclosed invention is the quantity which, when administered to a subject in need of treatment, improves the prognosis of the subject, e.g., delays the onset of and/or reduces the severity of one or more of the subject's symptoms associated with a fungal infection. The amount of the disclosed compound to be administered to a subject will depend on the particular disease, the mode of administration, and the characteristics of the subject, such as general health, other diseases, age, sex, genotype, body weight and tolerance to drugs. The skilled artisan will be able to determine appropriate dosages depending on these and other factors. Effective amounts of the disclosed compounds typically range between about 0.1mg/kg body weight per day and about 1000 mg/kg body weight per day, and preferably between 1 mg/kg body weight per day and 100 mg/kg body weight per day.

The present invention is directed to treating disorders mediated by pro-inflammatory cytokines such as, for example, TNF- α or HMGB-1. Non-limiting examples of disorders that can be treated using compounds, pharmaceutical compositions and methods of the present invention include appendicitis, peptic, gastric or duodenal ulcers, peritonitis, pancreatitis, ulcerative, pseudomembranous, acute or ischemic colitis, diverticulitis, epiglottitis, achalasia, cholangitis, cholecystitis, hepatitis, Crohn's disease, enteritis, Whipple's disease, asthma, allergy, anaphylactic shock, immune complex disease, organ ischemia, reperfusion injury, organ necrosis, hay fever, sepsis, septicemia, endotoxic shock, cachexia, hyperpyrexia, eosinophilic granuloma, granulomatosis, sarcoidosis, septic abortion, epididymitis, vaginitis, prostatitis, urethritis, bronchitis, emphysema, rhinitis, cystic fibrosis, pneumonitis, pneumoultramicroscopic silicovolcanoconiosis, alveolitis, bronchiolitis, pharyngitis, pleurisy, sinusitis, influenza, respiratory syncytial virus infection, herpes

infection, HIV infection, hepatitis B virus infection, hepatitis C virus infection, disseminated bacteremia, Dengue fever, candidiasis, malaria, filariasis, amebiasis, hydatid cysts, burns, dermatitis, dermatomyositis, sunburn, urticaria, warts, wheals, vasculitis, angiitis, endocarditis, arteritis, atherosclerosis, thrombophlebitis, pericarditis, myocarditis, myocardial ischemia, periarteritis nodosa, rheumatic fever, Alzheimer's disease, coeliac disease, congestive heart failure, adult respiratory distress syndrome, meningitis, encephalitis, multiple sclerosis, cerebral infarction, cerebral embolism, Guillame-Barre syndrome, neuritis, neuralgia, spinal cord injury, paralysis, uveitis, arthritides, arthralgias, osteomyelitis, fasciitis, Paget's disease, gout, periodontal disease, rheumatoid arthritis, synovitis, myasthenia gravis, thyroiditis, systemic lupus erythematosus, Goodpasture's syndrome, Behcets's syndrome, allograft rejection, graft-versus-host disease, Type I diabetes, ankylosing spondylitis, Berger's disease, Type I diabetes, ankylosing spondylitis, Retier's syndrome, or Hodgkins disease. In more preferred embodiments, the condition is appendicitis, peptic, gastric or duodenal ulcers, peritonitis, pancreatitis, ulcerative, pseudomembranous, acute or ischemic colitis, hepatitis, Crohn's disease, asthma, allergy, anaphylactic shock, organ ischemia, reperfusion injury, organ necrosis, hay fever, sepsis, septicemia, endotoxic shock, cachexia, septic abortion, disseminated bacteremia, burns, Alzheimer's disease, coeliac disease, congestive heart failure, adult respiratory distress syndrome, cerebral infarction, cerebral embolism, spinal cord injury, paralysis, allograft rejection and graft-versus-host disease.

Preferably, the disorder is peritonitis, pancreatitis, ulcerative colitis, Crohn's disease, asthma, organ ischemia, reperfusion ischemia, sepsis, cachexia, burns, myocardial ischemia, adult respiratory distress syndrome, multiple sclerosis, rheumatoid arthritis, systemic lupus erythematosus, chronic obstructive pulmonary disease, psoriasis, Behcet's syndrome, allograft rejection or graft-versus-host disease. Even more preferably, the disorder is sepsis.

The disclosed compounds can also be co-administered with one or more additional agents such as antibiotics, other anti-inflammatories (e.g. ibuprofen, prednisone (corticosteroid) or pentoxifylline), anti-fungals, (e.g. Amphotericin B, Fluconazole, Ketoconazol, and Itraconazol) steroids, decongestants, bronchodialators, and the like. The

disclosed compounds can also be co-administered with anti-TNF agents (e.g. infliximab, etanercept, adalimumab, CDP870, CDP571, Lenercept and Thalidomide). The formulation may also contain preserving agents, solubilizing agents, chemical buffers, surfactants, emulsifiers, colorants, odorants and sweeteners. The compounds of the present invention
5 may be co-administered with one or more additional agents separately or in the same formulation.

The excipient included with the compounds of the pharmaceutical compositions of the invention is chosen based on the expected route of administration of the composition in therapeutic applications. The route of administration of the composition depends on the
10 condition to be treated. For example, intravenous injection may be preferred for treatment of a systemic disorder such as endotoxic shock, and oral administration may be preferred to treat a gastrointestinal disorder such as a gastric ulcer. The route of administration and the dosage of the composition to be administered can be determined by the skilled artisan without undue experimentation in conjunction with standard dose-response studies. Relevant
15 circumstances to be considered in making those determinations include the condition or conditions to be treated, the choice of composition to be administered, the age, weight, and response of the individual patient, and the severity of the patient's symptoms. Thus, depending on the condition, the antibody composition can be administered orally, parenterally, intranasally, vaginally, rectally, lingually, sublingually, buccally, intrabuccally
20 and transdermally to the patient.

Accordingly, compositions designed for oral, lingual, sublingual, buccal and intrabuccal administration can be made without undue experimentation by means well known in the art, for example, with an inert diluent or with an edible carrier. The compositions may be enclosed in gelatin capsules or compressed into tablets. For the purpose of oral
25 therapeutic administration, the pharmaceutical compositions of the present invention may be incorporated with excipients and used in the form of tablets, troches, capsules, elixirs, suspensions, syrups, wafers, chewing gums and the like.

Tablets, pills, capsules, troches and the like may also contain binders, recipients, disintegrating agent, lubricants, sweetening agents, and flavoring agents. Some examples of
30 binders include microcrystalline cellulose, gum tragacanth or gelatin. Examples of

excipients include starch or lactose. Some examples of disintegrating agents include alginic acid, corn starch and the like. Examples of lubricants include magnesium stearate or potassium stearate. An example of a glidant is colloidal silicon dioxide. Some examples of sweetening agents include sucrose, saccharin and the like. Examples of flavoring agents include peppermint, methyl salicylate, orange flavoring and the like. Materials used in preparing these various compositions should be pharmaceutically pure and non-toxic in the amounts used.

Various other materials may be present as coatings or to modify the physical form of the dosage unit. For instance, tablets may be coated with shellac, sugar or both. A syrup or elixir may contain, in addition to the active ingredient, sucrose as a sweetening agent, methyl and propylparabens as preservatives, a dye and a flavoring such as cherry or orange flavor, and the like.

For vaginal administration, a pharmaceutical composition may be presented as pessaries, tampons, creams, gels, pastes, foams or spray.

The present invention includes nasally administering to the mammal a therapeutically effective amount of the composition. As used herein, nasally administering or nasal administration includes administering the composition to the mucous membranes of the nasal passage or nasal cavity of the patient. As used herein, pharmaceutical compositions for nasal administration of a composition include therapeutically effective amounts of the agonist prepared by well-known methods to be administered, for example, as a nasal spray, nasal drop, suspension, gel, ointment, cream or powder. Administration of the composition may also take place using a nasal tampon or nasal sponge.

For topical administration, suitable formulations may include biocompatible oil, wax, gel, powder, polymer, or other liquid or solid carriers. Such formulations may be administered by applying directly to affected tissues, for example, a liquid formulation to treat infection of conjunctival tissue can be administered dropwise to the subject's eye, or a cream formulation can be administer to a wound site.

The compositions of the present invention can be administered parenterally such as, for example, by intravenous, intramuscular, intrathecal or subcutaneous injection. Parenteral administration can be accomplished by incorporating the antibody compositions of the

present invention into a solution or suspension. Such solutions or suspensions may also include sterile diluents such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents. Parenteral formulations may also include antibacterial agents such as, for example, benzyl alcohol or methyl parabens, antioxidants such as, for example, ascorbic acid or sodium bisulfite and chelating agents such as EDTA. Buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose may also be added. The parenteral preparation can be enclosed in ampules, disposable syringes or multiple dose vials made of glass or plastic.

Rectal administration includes administering the pharmaceutical compositions into the rectum or large intestine. This can be accomplished using suppositories or enemas. Suppository formulations can easily be made by methods known in the art. For example, suppository formulations can be prepared by heating glycerin to about 120°C, dissolving the pharmaceutical composition in the glycerin, mixing the heated glycerin after which purified water may be added, and pouring the hot mixture into a suppository mold.

Transdermal administration includes percutaneous absorption of the composition through the skin. Transdermal formulations include patches, ointments, creams, gels, salves and the like.


In addition to the usual meaning of administering the formulations described herein to any part, tissue or organ whose primary function is gas exchange with the external environment, for purposes of the present invention, "pulmonary" will also mean to include a tissue or cavity that is contingent to the respiratory tract, in particular, the sinuses. For pulmonary administration, an aerosol formulation containing the active agent, a manual pump spray, nebulizer or pressurized metered-dose inhaler as well as dry powder formulations are contemplated. Suitable formulations of this type can also include other agents, such as antistatic agents, to maintain the disclosed compounds as effective aerosols.

A drug delivery device for delivering aerosols comprises a suitable aerosol canister with a metering valve containing a pharmaceutical aerosol formulation as described and an actuator housing adapted to hold the canister and allow for drug delivery. The canister in the drug delivery device has a head space representing greater than about 15% of the total

volume of the canister. Often, the polymer intended for pulmonary administration is dissolved, suspended or emulsified in a mixture of a solvent, surfactant and propellant. The mixture is maintained under pressure in a canister that has been sealed with a metering valve.

5 Synthesis of the Compounds of the Invention


The compounds of the present invention can be synthesized according to the following synthetic scheme:



- 10 Ar^a and Ar^b are each independently 6-member substituted or unsubstituted heteroaryl or aryl group as defined above (including variables Ar_1 , Ar_2 , Ar and ring **D**). This procedure is based on the method of isoxazoline synthesis found in Wityak *et al.*, J. Med. Chem. (1997) 40:50-60, incorporated herein by reference.

In particular, compounds (VI f), (VI d) and (XII a) can be prepared according to the
15 schemes (2), (3) and (4), respectively.


1) Compound (VI f):



- 20 A stirring solution of 3-pyridinecarboxaldehyde oxime (3.00 g, 24.6 mmol) and 4-vinylpyridine (8.0 mL, 75 mmol) in THF (60 mL) was chilled by an ice bath. To this solution, a 5% solution of NaOCl (95 mL) was added slowly by an addition funnel. When the addition was complete, the ice bath was removed and the reaction mixture was allowed to warm to room temperature. The reaction was quenched with 5% citric acid. Ether was added
25 to the reaction mixture was separated from the aqueous layer.

The aqueous layer was again extracted with ether, and the combined ether layers were dried with MgSO₄ and concentrated on the roto-evaporator. The crude product was loaded onto a silica column and eluted with 0, 10, 20, and 30% EtOAc/heptane. The isoxazoline came off the column in the 20-30% regime as a ~95% spectroscopically (¹H NMR) pure solid (1.32 g, 23.9%). The compound was crystallized from EtOAc/heptane.


2) Compound (VI d):



A stirring solution of 3-pyridinecarboxaldehyde oxime (3.00 g, 24.6 mmol) and 2-vinylpyridine (75 mmol) in THF (60 mL) was chilled by an ice bath. To this solution, a 5% solution of NaOCl (95 mL) was added slowly by an addition funnel. When the addition was complete, the ice bath was removed and the reaction mixture was allowed to warm to room temperature. The reaction was quenched with 5% citric acid. Ether was added to the reaction mixture and was separated from the aqueous layer.

The aqueous layer was again extracted with ether, and the combined ether layers were dried with MgSO₄ and concentrated on the roto-evaporator. The crude product was loaded onto a silica column and eluted with 0, 10, 20, and 30% EtOAc/heptane. The isoxazoline came off the column in the 20-30% regime as a ~95% spectroscopically (¹H NMR) pure solid (1.53 g, 27.7%). The compound was crystallized from EtOAc/heptane.

3) Compound (XII a):



A stirring solution of 4-pyridinecarboxaldehyde oxime (3.00 g, 24.6 mmol) and Styrene (75 mmol) in THF (60 mL) was chilled by an ice bath. To this solution, a 5% solution of NaOCl (95 mL) was added slowly by an addition funnel. When the addition was complete, the ice bath was removed and the reaction mixture was allowed to warm to room temperature. The reaction was quenched with 5% citric acid. Ether was added to the reaction mixture and was separated from the aqueous layer.

The aqueous layer was again extracted with ether, and the combined ether layers were dried with MgSO₄ and concentrated on the roto-evaporator. The crude product was loaded onto a silica column and eluted with 0, 10, 20, and 30% EtOAc/heptane. The isoxazoline came off the column in the 10-20 % regime as a > 95% spectroscopically (¹H NMR) pure solid (2.1 g, 38%). The compound was crystallized from EtOAc/heptane.

Other compounds of the invention can be prepared with these methods using the appropriate starting materials. In some instances, where the starting materials have substituent groups, it may be necessary to use protecting groups. Suitable protecting groups are known to the one of ordinary skill in the art and are found, for example, in Green and Wuts, "Protective Groups in Organic Synthesis", John Wiley and Sons (1991), the entire relevant teaching of which are incorporated herein by reference.

EXEMPLIFICATION

The practice of the present invention will employ, unless otherwise indicated, conventional techniques of cell culture, molecular biology, microbiology, cell biology, and immunology, which are well within the skill of the art. Such techniques are fully explained in the literature. See, e.g., Sambrook et al., 1989, "Molecular Cloning: A Laboratory Manual", Cold Spring Harbor Laboratory Press; Ausubel et al. (1995), "Short Protocols in Molecular Biology", John Wiley and Sons; Methods in Enzymology (several volumes); Methods in Cell Biology (several volumes), and Methods in Molecular Biology (several volumes).

Example 1 Compound of Formula (VI) Suppresses TNF Release from LPS-Stimulated RAW Cells in a Dose-Dependent Manner

Experiments were performed to determine the effect of compound (VI f) on the proinflammatory activity of TNF α . Murine RAW 264.7 macrophage-like cells (American Type Tissue Culture Collection, Rockville, Md., USA) were grown under DMEM supplemented with 10% fetal bovine serum, penicillin and streptomycin. The cells were seeded in 24-well tissue culture plates in Opti-MEM 1 medium and used at 90% confluence. The cells were treated with either compound (VI f) or nicotine (Sigma) at 0, 0.001, 0.01, 0.1, 1.0, 10 or 100 μ M. Five minutes after the addition of compound (VI f) or nicotine, the cells were treated with LPS (500 ng/ml). Supernatants were collected after 4 hours and TNF concentration was measured by mouse ELISA kit (R&D Systems Inc., Minneapolis, MN).

The results are shown in FIG. 1, which clearly shows that compound (VI f) inhibited TNF α production in a dose-dependent manner. The efficacy of compound (VI f) is comparable to or better than that of the nicotine control.

Example 2 Compound of Formula (VI f) Suppresses HMGB-1 Release from LPS-Stimulated RAW Cells in a Dose-Dependent Manner

Experiments were performed to determine the effect of compound (VI f) on the proinflammatory effect of HMGB-1. Murine RAW 264.7 macrophage-like cells were grown under DMEM supplemented with 10% fetal bovine serum and 1% glutamine. When the cells were 70-80% confluent, the medium was replaced by serum-free Opti-MEM I medium. Compound (VI f) was added at 0, 0.01, 0.1, 1 or 10 μ M or nicotine was added at 0, 0.1, 1, 10 and 100 μ M. Five minutes after the addition of compound (VI f) or nicotine, the cultures were treated with LPS (500 ng/ml). Culture medium was collected after 24 hours. The culture medium was concentrated with a CentriconTM 10 filter, then analyzed by western blot using anti-HMGB1 polyclonal antisera and standard methods. The results are presented as percent stimulation of LPS treatment alone.

The results are shown in FIG. 2, which clearly shows that compound (VI f) inhibited HMGB1 production in a dose-dependent manner. The efficacy of compound (VI f) is comparable to or better than that of nicotine control.

Example 3 Compounds of the Invention Exhibit Low IC₅₀ for TNF α Release

Experiments were performed to determine the concentration of several compounds of the present invention at which TNF α release was inhibited by 50 % (IC₅₀).

- Using murine RAW 264.7 macrophage-like cells and the protocol described in
- 5 Example 1, IC₅₀ of compounds (VI f), (VI b), (XII b), (VI e), (VI g) and (XII a) have been measured. The results are summarized in FIG. 3. As can be seen from FIG. 3, four out of six compounds tested exhibit IC₅₀ values comparable or better than nicotine.

Example 4 Compound (VI f) Protects Mice from Septic Shock in CLP Model

- 10 Cecal Ligation and Puncture (CLP) was performed as described in Fink and Heard, J. of Surg. Res. 49:186-196 (1990), Wichman *et al.*, Crit. Care Med. 26:2078-2086 (1998) and Remick *et al.*, Shock 4:89-95 (1995). Briefly, Balb/c mice were anesthetized with 75 mg/kg Ketamine (Fort Dodge, Fort Dodge, Iowa) and 20 mg/kg of xylazine (Bohringer Ingelheim, St. Joseph, MO) intramuscularly. A midline incision was performed, and the cecum was
- 15 isolated. A 6-0 prolene suture ligature was placed at a level 5.0 mm from the cecal tip away from the ileocecal valve.

- The ligated cecal stump was then punctured once with a 22-gauge needle, without direct extrusion of stool. The cecum was then placed back into its normal intra-abdominal position. The abdomen was then closed with a running suture of 6-0 prolene in two layers,
- 20 peritoneum and fascia separately to prevent leakage of fluid. All animals were resuscitated with a normal saline solution administered sub-cutaneously at 20 ml/kg of body weight. Each mouse received a subcutaneous injection of imipenem (0.5 mg/mouse) (Primaxin, Merck & Co., Inc., West Point, PA) 30 minutes after the surgery. Animals were then allowed to recuperate. Mortality was recorded for up to 1 week after the procedure;
- 25 survivors were followed for 2 weeks to ensure no late mortalities had occurred.

- The results are presented in FIG. 4, which shows the number of surviving animals following treatment either by compound (VI f) or an "empty" carrier solution. As can be seen, only two mice died in the compound (VI f)-treated group as compared with six in the untreated control. By day 5, only one treated mouse died compared to six in the control
- 30 group. These results demonstrate that compound VI f improved survival of septic mice.

Example 5 Compounds (VI d) and (VI f) Inhibit TNFa Production in LPS-stimulated Macrophages

Human macrophage cultures were prepared as follows. Buffy coats were collected from the blood of healthy individual donors to the Long Island Blood Bank Services (Melville, N.Y.). Primary blood mononuclear cells were isolated by density-gradient centrifugation through Ficoll/Hypaque (Pharmacia, N.J.), suspended (8×10^6 cells/ml) in RPMI 1640 medium supplemented with 10% heat inactivated human serum (Gemini Bio-Products, Inc., Calabasas, Calif.), and seeded in flasks (PRIMARIA; Beckton and Dickinson Labware, Franklin Lakes, N.J.). After incubation for 2 hours at 37 °C, adherent cells were washed extensively, treated briefly with 10 mM EDTA, detached, resuspended (10^6 cells/ml) in RPMI medium (10% human serum), supplemented with human macrophage colony stimulating factor (MCSF; Sigma Chemical Co., St. Louis, Mo.; 2 ng/ml), and seeded onto 24-well tissue culture plates (PRIMARIA; Falcon) (10×10^6 cells/well). Cells were allowed to differentiate for 7 days in the presence of MCSF. On day 7 the cells were washed 3 times with 1x Dulbecco's phosphate buffered saline (PBS, GibcoBRL, Life Technologies, Rockville, Md.), fresh medium devoid of MCSF was added, and experiments performed as indicated.

Primary human macrophage cultures were established by incubating human peripheral blood mononuclear cells in the presence of macrophage colony stimulating factor (MCSF; Sigma Chemical Co., St. Louis, Mo.). These cells were used in experiments to determine the effects of the compounds of the present invention on TNF levels in macrophage cultures conditioned by exposure to LPS for 4 hours. In those experiments, compounds (VI d) and (VI f) as well as control (nicotine; Sigma Chemical Co., St. Louis, Mo.) was added to human macrophage cultures at the indicated concentrations (FIG. 5). LPS was added five minutes later (100 ng/ml), and conditioned supernatants collected after 4 hours of stimulation for subsequent analysis by TNF enzyme-linked immunosorbent assay (ELISA). All the experimental conditions were performed in triplicate. Data from nine separate macrophage preparations are shown as Mean \pm SEM; n=9.

As can be seen from FIG. 5, compounds (VI d) and (VI f) inhibited TNF release in a dose dependent manner with efficacy comparable to or better than that of the control.

Example 6 α -Bungarotoxin Overrides Inhibitory Effect of Compound (VI f) on TNF α

5 Release

Experiments were performed to determine whether the inhibitory effect of the compound (VI f) on proinflammatory cytokine TNF α can be overridden by addition of α -bungarotoxin, an antagonist that binds to a subset of cholinergic receptors including the nicotinic $\alpha 7$ receptor (Lindstrom, 1995; Leonard et al, 2001).

10 Murine RAW 264.7 macrophage-like cells were subjected to the protocol as described in Example 1. The cells were treated with α -bungarotoxin at the indicated concentration before treatment with compound (VI f).

As shown in FIG. 6, addition of α -bungarotoxin partially abrogated the inhibitory effect of compound (VI f) on LPS-mediated TNF α release, which demonstrates that this inhibition is at least partially dependent on an α -bungarotoxin sensitive receptor on the RAW 264.7 cells.


15 While this invention has been particularly shown and described with references to preferred embodiments thereof, it will be understood by those skilled in the art that various changes in form and details may be made therein without departing from the scope of the invention encompassed by the appended claims.

20

CLAIMS

What is claimed is:

- 5 1. A compound of Formula (I) or a pharmaceutically acceptable salt thereof:



wherein:

- 10 Ar_1 and Ar_2 are independently a monocyclic six-member optionally substituted heteroaryl group;


A_1 is =N- or -NR^a- and A_2 is O or S;

R^a is H or C1-C6 alkyl;

- 15 R_1 is selected from -H, C1-C6 alkyl, phenyl, C1-C6 haloalkyl, halogen, -OH, -OR^b, C1-C6 hydroxyalkyl, C1-C6 alkoxyalkyl, -O(C1-C6 haloalkyl), -SH, -SR^b, -NO₂, -CN, -NR^bCO₂ R^b, -NR^bC(O) R^b, -CO₂R^b, -C(O)R^b, -C(O)N(R^b)₂, -OC(O)R^b and -NR^bR^b; and

each R^b is H or a C1-C6 alkyl group.

- 20 2. The compound of Claim 1 represented by Formula (II):




wherein:

B_1 through B_5 and D_1 through D_5 are independently N or CR^c ,
provided that from one to three of B_1 through B_5 and from one to three of D_1
through D_5 are N;


each R^c is independently -H, C1-C6 alkyl, halogen, C1-C6 haloalkyl,
- R^o , -OH, -OR o , -O(C1-C6 haloalkyl), -SH, -SR o , -NO $_2$, -CN, -NHCO $_2R^o$,
-NHC(O)H, -NHC(O) R^o , -CO $_2H$, -CO $_2R^o$, -C(O)H, -C(O) R^o , -C(O)NHR o ,
-OC(O) R^o , -S(O) $_2R^o$, -SO $_2NH_2$, -S(O) R^o , -NHSO $_2R^o$, -N(R^o) $_2$, -NR o C(O) R^o ,
or -C(O)N(R^o) $_2$ or a C1-C6 alkyl group substituted with R^o , -OH, -OR o , -SH,
-SR o , -NO $_2$, -CN, -NHCO $_2R^o$, -NHC(O)H, -NHC(O) R^o , -CO $_2H$, -CO $_2R^o$,
-C(O)H, -C(O) R^o , -C(O)NHR o , -OC(O) R^o , -S(O) $_2R^o$, -SO $_2NH_2$, -S(O) R^o ,
-NHSO $_2R^o$; and

R^o is independently, C1-C6 alkyl, aryl or heteroaryl group and wherein
the aryl or heteroaryl group represented by R^o is optionally substituted with
one or more halogen, methyl or methoxy groups.

3. The compound of Claim 2 wherein the compound is represented by Formula (III):





4. The compound of Claim 3 wherein the compound is represented by Formula (IV):



wherein one of B₁ through B₃ and one of D₁ through D₃ are N.

5. The compound of Claim 4 wherein R₁ is -H or a C1-C3 alkyl optionally substituted with a halogen or a hydroxyl.
6. The compound of Claim 5 wherein
 R^c is -H, halogen, -NO₂, -CN, C1-C3 alkyl, C1-C3 haloalkyl, C1-C3 hydroxyalkyl, C1-C3 alkoxyalkyl, -N(R^d)₂, -NR^dC(O)R^d, or -C(O)N(R^d)₂; and
 each R^d is H or a C1-C3 alkyl group.
7. The compound of Claim 6 selected from the group consisting of:







5 wherein:


R' and R'' are independently -H, halogen, -NO₂, -CN, C1-C3 alkyl, C1-C3 haloalkyl, C1-C3 hydroxyalkyl, C1-C3 alkoxyalkyl, -N(R^d)₂, -NR^dC(O)R^d, or -C(O)N(R^d)₂.

10 8. The compound of Claim 7 selected from the group consisting of:






9. The compound of Claim 8 represented by Formula (VI d):




5

10. The compound of Claim 8 represented by Formula (VI f):



- 10 11. A pharmaceutical composition comprising a pharmaceutically acceptable carrier or diluent and a compound represented by Formula (I a) or a pharmaceutically acceptable salt thereof:



15

wherein:


Ar_1 and Ar_2 are independently a monocyclic six-member optionally substituted heteroaryl group;

A_1 is $=N-$ or $-NR^a-$ and A_2 is O or S;

R^a is H or C1-C6 alkyl;

R_1 is selected from -H, C1-C6 alkyl, phenyl, C1-C6 haloalkyl, halogen, -OH, -OR^b, C1-C6 hydroxyalkyl, C1-C6 alkoxyalkyl, -O(C1-C6 haloalkyl), -SH, -SR^b, -NO₂, -CN, -NR^bCO₂R^b, -NR^bC(O)R^b, -CO₂R^b, -C(O)R^b, -C(O)N(R^b)₂, -OC(O)R^b and -NR^bR^b; and each R^b is H or a C1-C6 alkyl group.

12. The composition of Claim 11 wherein the compound is represented by Formula (II a):




wherein:

B_1 through B_5 and D_1 through D_5 are independently N or CR^c, provided that from one to three of B_1 through B_5 and from one to three of D_1 through D_5 are N;

each R^c is independently -H, C1-C6 alkyl, halogen, C1-C6 haloalkyl, -R^o, -OH, -OR^o, -O(C1-C6 haloalkyl), -SH, -SR^o, -NO₂, -CN, -NHCO₂R^o, -NHC(O)H, -NHC(O)R^o, -CO₂H, -CO₂R^o, -C(O)H, -C(O)R^o, -C(O)NHR^o, -OC(O)R^o, -S(O)₂R^o, -SO₂NH₂, -S(O)R^o, -NHCO₂R^o, -N(R^o)₂, -NR^oC(O)R^o, -C(O)N(R^o)₂ or a C1-C6 alkyl group substituted with R^o, -OH, -OR^o, -SH, -SR^o, -NO₂, -CN, -NHCO₂R^o, -NHC(O)H, -NHC(O)R^o, -CO₂H, -CO₂R^o, -C(O)H, -C(O)R^o, -C(O)NHR^o, -OC(O)R^o, -S(O)₂R^o, -SO₂NH₂, -S(O)R^o or -NHCO₂R^o;


R^0 is independently C1-C6 alkyl, aryl or heteroaryl group and wherein the aryl or heteroaryl group represented by R^0 is optionally substituted with one or more halogen, methyl or methoxy groups.

- 5 13. The composition of Claim 12 wherein the compound is represented by Formula (III):



14. The composition of Claim 13 wherein the compound is represented by Formula (IV):


10



wherein one of B_1 through B_3 and one of D_1 through D_3 are N.

- 15 15. The composition of Claim 14 wherein R_1 is -H or a C1-C3 alkyl optionally substituted with a halogen or a hydroxyl.
16. The composition of Claim 15 wherein
- R^c is -H, halogen, -NO₂, -CN, C1-C3 alkyl, C1-C3 haloalkyl, C1-C3 hydroxyalkyl or C1-C3 alkoxyalkyl, -N(R^d)₂, -NR^dC(O)R^d, or -C(O)N(R^d)₂; and
- 20 each R^d is H or a C1-C3 alkyl group.


17. The composition of Claim 16 wherein the compound is selected from the group consisting of:




10 wherein

R' and R'' are independently -H, halogen, $-NO_2$, $-CN$, C1-C3 alkyl, C1-C3 haloalkyl, C1-C3 hydroxyalkyl, C1-C3 alkoxyalkyl, $-N(R^d)_2$, $-NR^dC(O)R^d$, or $-C(O)N(R^d)_2$.


- 15 18. The composition of Claim 17 wherein the compound is selected from the group consisting of:






5


19. The composition of Claim 18 wherein the compound is represented by Formula (VI d):



- 10 20. The composition of Claim 18 wherein the compound is represented by Formula (VI f):



21. A pharmaceutical composition comprising a pharmaceutically acceptable carrier or diluent and a compound represented by Formula (VII) or a pharmaceutically acceptable salt thereof:



wherein:

Ar is an optionally substituted, monocyclic, six-member heteroaryl;

A₁ is =N- or -NR^a- and A₂ is O or S;

10 R₁ is -H, C1-C6 alkyl, phenyl, C1-C6 haloalkyl, halogen, -OH, -OR^b, C1-C6 hydroxyalkyl, C1-C6 alkoxyalkyl, -O(C1-C6 haloalkyl), -SH, -SR^b, -NO₂, -CN, -NR^bCO₂ R^b, -NR^bC(O) R^b, -CO₂R^b, -C(O)R^b, -C(O)N(R^b)₂, -OC(O)R^b or -NR^bR^b;

each R^a is -H or C1-C6 alkyl;

15 each R^b is -H or a C1-C6 alkyl group; and

ring D is optionally substituted with zero, one or more substituents other than amide and is not an alkylphenol.

22. The pharmaceutical composition of Claim 21 wherein:

20 ring D is optionally substituted with zero, one or more substituents represented by R₂;


each R₂ is independently -H, halogen, C1-C6 haloalkyl, -R^o, -OH, -OR^o, -O(C1-C6 haloalkyl), -SH, -SR^o, -NO₂, -CN, -NHCO₂R^o, -NHC(O)H, -NHC(O)R^o, -CO₂H, -CO₂R^o, -C(O)H, -C(O)R^o, -OC(O)R^o, -S(O)₂R^o, -SO₂NH₂, -S(O)R^o, -NHSO₂R^o, or a C1-C6 alkyl group substituted with R^o, -OH, -OR^o, -SH, -SR^o, -NO₂,

25

-CN, -NHCO₂R^o, -NHC(O)H, -NHC(O)R^o, -CO₂H, -CO₂R^o, -C(O)H, -C(O)R^o,
-C(O)NHR^o, -OC(O)R^o, -S(O)₂R^o, -SO₂NH₂, -S(O)R^o or -NHSO₂R^o;

R^o is independently -H, C1-C6 alkyl, aryl or heteroaryl group wherein the aryl
or heteroaryl group represented by R^o is optionally substituted with one or more
halogen, methyl or methoxy groups.

23. The composition of Claim 22 wherein the compound is represented by Formula
(VIII):




wherein:

B₁ through B₅ are independently N or CR^c, provided that from one to
three of B₁ through B₅ are N;


each R^c is independently -H, C1-C6 alkyl, halogen, C1-C6 haloalkyl,
-R^o, -OH, -OR^o, -O(C1-C6 haloalkyl), -SH, -SR^o, -NO₂, -CN, -NHCO₂R^o,
-NHC(O)H, -NHC(O)R^o, -CO₂H, -CO₂R^o, -C(O)H, -C(O)R^o, -C(O)NHR^o,
-OC(O)R^o, -S(O)₂R^o, -SO₂NH₂, -S(O)R^o, -NHSO₂R^o, -N(R^o)₂, -NR^oC(O)R^o,
-C(O)N(R^o)₂ or a C1-C6 alkyl group substituted with R^o, -OH, -OR^o, -SH,
-SR^o, -NO₂, -CN, -NHCO₂R^o, -NHC(O)H, -NHC(O)R^o, -CO₂H, -CO₂R^o,
-C(O)H, -C(O)R^o, -C(O)NHR^o, -OC(O)R^o, -S(O)₂R^o, -SO₂NH₂, -S(O)R^o,
-NHSO₂R^o.

24. The pharmaceutical composition of Claim 23 wherein the compound is represented
by formula (IX):

- 37 -



25. The composition of Claim 24 wherein the compound is represented by formula (X):



wherein:

one of B₁ through B₅ is N;

n is 0, 1 or 2; and

m is 0, 1 or 2.

26. The composition of Claim 25 wherein R₁ is -H or a C1-C3 alkyl optionally substituted with a halogen or a hydroxyl.


27. The composition of Claim 26 wherein:

each R₂ is independently -H, halogen, -NO₂, -CN, C1-C3 alkyl, C1-C3 haloalkyl, C1-C3 hydroxyalkyl or C1-C3 alkoxyalkyl;


each R^c is independently -H, halogen, -NO₂, -CN, C1-C3 alkyl, C1-C3 haloalkyl, C1-C3 hydroxyalkyl, C1-C3 alkoxyalkyl, -N(R^d)₂, -NR^dC(O)R^d, -C(O)N(R^d)₂; and

each R^d is H or a C1-C3 alkyl group.


28. The composition of Claim 27 wherein the compound is selected from the group of:



29. The pharmaceutical composition of Claim 28 wherein the compound is selected from the group consisting of:



30. A method of treating a subject with an inflammatory cytokine mediated disorder comprising administering to the subject a therapeutically effective amount of a compound represented by Formula (I) or a pharmaceutically acceptable salt thereof:



wherein:

Ar₁ and Ar₂ are independently monocyclic six-member optionally substituted heteroaryl groups;


A_1 is $=N-$ or $-NR^a-$ and A_2 is O or S;

R^a is -H or C1-C6 alkyl;

R_1 is y selected from -H, C1-C6 alkyl, phenyl, C1-C6 haloalkyl, halogen, -OH, $-OR^b$, C1-C6 hydroxyalkyl, C1-C6 alkoxyalkyl, $-O(C1-C6$ haloalkyl), -SH, $-SR^b$, $-NO_2$, -CN, $-NR^bCO_2R^b$, $-NR^bC(O)R^b$, $-CO_2R^b$, $-C(O)R^b$, $-C(O)N(R^b)_2$, $-OC(O)R^b$ and $-NR^bR^b$; and

R^b is -H or a C1-C6 alkyl group.

31. The method of Claim 30, wherein the compound is represented by Formula (II a):




wherein:

B_1 through B_5 and D_1 through D_5 are independently N or CR^c ,
provided that from one to three of B_1 through B_5 and from one to three of D_1 through D_5 are N;

each R^c is independently -H, C1-C6 alkyl, halogen, C1-C6 haloalkyl, $-R^o$, -OH, $-OR^o$, $-O(C1-C6$ haloalkyl), -SH, $-SR^o$, $-NO_2$, -CN, $-NHCO_2R^o$, $-NHC(O)H$, $-NHC(O)R^o$, $-CO_2H$, $-CO_2R^o$, $-C(O)H$, $-C(O)R^o$, $-C(O)NHR^o$, $-OC(O)R^o$, $-S(O)_2R^o$, $-SO_2NH_2$, $-S(O)R^o$, $-NHSO_2R^o$, $-N(R^o)_2$, $-NR^oC(O)R^o$, $-C(O)N(R^o)_2$ or a C1-C6 alkyl group substituted with R^o , -H, $-OR^o$, -SH, $-SR^o$, $-NO_2$, -CN, $-NHCO_2R^o$, $-NHC(O)H$, $-NHC(O)R^o$, $-CO_2H$, $-CO_2R^o$, $-C(O)H$, $-C(O)R^o$, $-C(O)NHR^o$, $-OC(O)R^o$, $-S(O)_2R^o$, $-SO_2NH_2$, $-S(O)R^o$ or $-NHSO_2R^o$; and


R^o is independently C1-C6 alkyl, aryl or heteroaryl group and wherein the aryl or heteroaryl group represented by R^o is optionally substituted with one or more halogen, methyl or methoxy groups.

- 5 32. The method of Claim 31 wherein the compound is represented by Formula (III):



33. The method of Claim 32 wherein the compound is represented by Formula (IV):

10












wherein one of B_1 through B_3 and one of D_1 through D_3 are N.

- 15 34. The method of Claim 33 wherein R_1 is -H or a C1-C3 alkyl, optionally substituted with a halogen or a hydroxyl.

35. The method of Claim 34 wherein

20 R^c is -H, halogen, -NO₂, -CN, C1-C3 alkyl, C1-C3 haloalkyl, C1-C3 hydroxyalkyl, C1-C3 alkoxyalkyl, -N(R^d)₂, -NR^dC(O)R^d, -C(O)N(R^d)₂; and each R^d is H or a C1-C3 alkyl group.



36. The method of Claim 35 wherein the compound is selected from the group consisting of:




10 wherein

R' and R'' are independently -H, halogen, -NO₂, -CN, C1-C3 alkyl, C1-C3 haloalkyl, C1-C3 hydroxyalkyl, C1-C3 alkoxyalkyl, -N(R^d)₂, -NR^dC(O)R^d, -C(O)N(R^d)₂.


- 15 37. The method of Claim 36 wherein the compound is selected from the group consisting of:






5

38. The method of Claim 37 wherein the compound is represented by Formula (VI d):



- 10 39. The method of Claim 37 wherein the compound is represented by Formula (VI f):




40. The method of Claim 30 wherein the inflammatory cytokine is TNF- α or HMGB-1.

41. The method of Claim 40 wherein the disorder is selected from the group consisting of
 appendicitis, peptic, gastric or duodenal ulcers, peritonitis, pancreatitis, ulcerative,
 pseudomembranous, acute or ischemic colitis, diverticulitis, epiglottitis, achalasia,
 5 cholangitis, cholecystitis, hepatitis, Crohn's disease, enteritis, Whipple's disease,
 asthma, allergy, anaphylactic shock, immune complex disease, organ ischemia,
 reperfusion injury, organ necrosis, hay fever, sepsis, septicemia, endotoxic shock,
 cachexia, hyperpyrexia, eosinophilic granuloma, granulomatosis, sarcoidosis, septic
 abortion, epididymitis, vaginitis, prostatitis, urethritis, bronchitis, emphysema,
 10 rhinitis, cystic fibrosis, pneumonitis, pneumoultramicroscopic silicovolcanoconiosis,
 alveolitis, bronchiolitis, pharyngitis, pleurisy, sinusitis, influenza, respiratory
 syncytial virus infection, herpes infection, HIV infection, hepatitis B virus infection,
 hepatitis C virus infection, disseminated bacteremia, Dengue fever, candidiasis,
 malaria, filariasis, amebiasis, hydatid cysts, burns, dermatitis, dermatomyositis,
 15 sunburn, urticaria, warts, wheals, vasculitis, angitis, endocarditis, arteritis,
 atherosclerosis, thrombophlebitis, pericarditis, myocarditis, myocardial ischemia,
 periarteritis nodosa, rheumatic fever, Alzheimer's disease, coeliac disease, congestive
 heart failure, adult respiratory distress syndrome, meningitis, encephalitis, multiple
 sclerosis, cerebral infarction, cerebral embolism, Guillame-Barre syndrome, neuritis,
 20 neuralgia, spinal cord injury, paralysis, uveitis, arthritides, arthralgias, osteomyelitis,
 fasciitis, Paget's disease, gout, periodontal disease, rheumatoid arthritis, synovitis,
 myasthenia gravis, thyroiditis, systemic lupus erythematosus, Goodpasture's
 syndrome, Behcets's syndrome, allograft rejection, graft-versus-host disease, Type I
 diabetes, ankylosing spondylitis, Berger's disease, Type I diabetes, ankylosing
 25 spondylitis, Retier's syndrome, or Hodgkins disease. In more preferred
 embodiments, the condition is appendicitis, peptic, gastric or duodenal ulcers,
 peritonitis, pancreatitis, ulcerative, pseudomembranous, acute or ischemic colitis,
 hepatitis, Crohn's disease, asthma, allergy, anaphylactic shock, organ ischemia,
 reperfusion injury, organ necrosis, hay fever, sepsis, septicemia, endotoxic shock,
 30 cachexia, septic abortion, disseminated bacteremia, burns, Alzheimer's disease,

coeliac disease, congestive heart failure, adult respiratory distress syndrome, cerebral infarction, cerebral embolism, spinal cord injury, paralysis, allograft rejection and graft-versus-host disease.

- 5 42. The method of Claim 41 wherein the disorder is selected from the group consisting of peritonitis, pancreatitis, ulcerative colitis, Crohn's disease, asthma, organ ischemia, reperfusion ischemia, sepsis, cachexia, burns, myocardial ischemia, adult respiratory distress syndrome, multiple sclerosis, rheumatoid arthritis, systemic lupus erythematosus, chronic obstructive pulmonary disease, psoriasis, Behcet's syndrome, allograft rejection and graft-versus-host disease.
- 10
43. The method of Claim 42 wherein the disorder is sepsis.
44. A method of treating a subject with an inflammatory cytokine mediated disorder comprising administering to the subject a therapeutically effective amount of a compound represented by Formula (VIIa) or a pharmaceutically acceptable salt thereof:
- 15



20 wherein

Ar is an optionally substituted, monocyclic, six-member heteroaryl;

A₁ is =N- or -NR^a- and A₂ is O or S;

R₁ is -H, C1-C6 alkyl, phenyl, C1-C6 haloalkyl, halogen, -OH, -OR^b, C1-C6 hydroxyalkyl, C1-C6 alkoxyalkyl, -O(C1-C6 haloalkyl), -SH, -SR^b, -NO₂, -CN, -NR^bCO₂ R^b, -NR^bC(O) R^b, -CO₂R^b, -C(O)R^b, -C(O)N(R^b)₂, -OC(O)R^b or -NR^bR^b;

25

each R^a is -H or C1-C6 alkyl group;

each R^b is -H or a C1-C6 alkyl group;

ring **D** is optionally substituted with zero, one or more substituents other than amide and is not an alkylphenol.

5

45. The pharmaceutical composition of Claim 44 wherein

ring **D** is optionally substituted with zero, one or more substituents represented by R_2 ;


each R_2 is independently -H, a halogen, C1-C6 haloalkyl, $-R^o$, $-OH$, $-OR^o$,
 10 $-O(C1-C6 \text{ haloalkyl})$, $-SH$, $-SR^o$, $-NO_2$, $-CN$, $-NHCO_2R^o$, $-NHC(O)H$, $-NHC(O)R^o$,
 $-CO_2H$, $-CO_2R^o$, $-C(O)H$, $-C(O)R^o$, $-OC(O)R^o$, $-S(O)_2R^o$, $-SO_2NH_2$, $-S(O)R^o$, $-$
 $NHSO_2R^o$, or a C1-C6 alkyl group substituted with R^o , $-OH$, $-OR^o$, $-SH$, $-SR^o$, $-NO_2$,
 $-CN$, $-NHCO_2R^o$, $-NHC(O)H$, $-NHC(O)R^o$, $-CO_2H$, $-CO_2R^o$, $-C(O)H$, $-C(O)R^o$,
 $-C(O)NHR^o$, $-OC(O)R^o$, $-S(O)_2R^o$, $-SO_2NH_2$, $-S(O)R^o$ or $-NHSO_2R^o$, and

15

R^o is independently -H, C1-C6 alkyl, aryl or heteroaryl group wherein the aryl or heteroaryl group represented by R^o is optionally substituted with one or more halogen, methyl or methoxy groups.

46. The method of Claim 45 wherein the compound is represented by Formula (VIIIa):

20




wherein:

B_1 through B_5 are independently N or CR^c , provided that from one to
 25 three of B_1 through B_5 are N;


each R° are independently -H, halogen, C1-C6 haloalkyl, $-R^{\circ}$, -OH, $-OR^{\circ}$, $-O(C1-C6 \text{ haloalkyl})$, -SH, $-SR^{\circ}$, $-NO_2$, -CN, $-NHCO_2R^{\circ}$, $-NHC(O)H$, $-NHC(O)R^{\circ}$, $-CO_2H$, $-CO_2R^{\circ}$, $-C(O)H$, $-C(O)R^{\circ}$, $-C(O)NHR^{\circ}$, $-OC(O)R^{\circ}$, $-S(O)_2R^{\circ}$, $-SO_2NH_2$, $-S(O)R^{\circ}$, $-NHSO_2R^{\circ}$, $-N(R^{\circ})_2$, $-NR^{\circ}C(O)R^{\circ}$, $-C(O)N(R^{\circ})_2$ or a C1-C6 alkyl group substituted with R° , -OH, $-OR^{\circ}$, -SH, $-SR^{\circ}$, $-NO_2$, -CN, $-NHCO_2R^{\circ}$, $-NHC(O)H$, $-NHC(O)R^{\circ}$, $-CO_2H$, $-CO_2R^{\circ}$, $-C(O)H$, $-C(O)R^{\circ}$, $-C(O)NHR^{\circ}$, $-OC(O)R^{\circ}$, $-S(O)_2R^{\circ}$, $-SO_2NH_2$, $-S(O)R^{\circ}$ or $-NHSO_2R^{\circ}$; and

R° is independently -H, C1-C6 alkyl, aryl or heteroaryl group wherein the aryl or heteroaryl group represented by R° is optionally substituted with one or more halogen, methyl or methoxy groups.

47. The method of Claim 46 wherein the compound is represented by formula (IX):



48. The method of Claim 47 wherein the compound is represented by formula (X):



wherein

one of B1 through B3 is N;

n is 0, 1 or 2; and

m is 0, 1 or 2.

49. The composition of Claim 48 wherein R_1 is -H or a C1-C3 alkyl optionally substituted with a halogen or a hydroxyl.

5



50. The method of Claim 49 wherein:

each R^c and each R_2 are independently -H, halogen, $-\text{NO}_2$, $-\text{CN}$, C1-C3 alkyl, C1-C3 haloalkyl, C1-C3 hydroxyalkyl, C1-C3 alkoxyalkyl, $-\text{N}(\text{R}^d)_2$, $-\text{NR}^d\text{C}(\text{O})\text{R}^d$, $-\text{C}(\text{O})\text{N}(\text{R}^d)_2$; and

10


each R^d is H or a C1-C3 alkyl group.

51. The method of Claim 50 wherein the compound is selected from the group of:



15

52. The method of Claim 51 wherein the compound is selected from the group consisting of:



20

53. The method of Claim 44 wherein the inflammatory cytokine is TNF- α or HMGB-1.

54. The method of Claim 53 wherein the disorder is selected from the group consisting of
 5 appendicitis, peptic, gastric or duodenal ulcers, peritonitis, pancreatitis, ulcerative,
 pseudomembranous, acute or ischemic colitis, diverticulitis, epiglottitis, achalasia,
 cholangitis, cholecystitis, hepatitis, Crohn's disease, enteritis, Whipple's disease,
 asthma, allergy, anaphylactic shock, immune complex disease, organ ischemia,
 reperfusion injury, organ necrosis, hay fever, sepsis, septicemia, endotoxic shock,
 cachexia, hyperpyrexia, eosinophilic granuloma, granulomatosis, sarcoidosis, septic
 10 abortion, epididymitis, vaginitis, prostatitis, urethritis, bronchitis, emphysema,
 rhinitis, cystic fibrosis, pneumonitis, pneumoultramicroscopic silicovolcanoconiosis,
 alveolitis, bronchiolitis, pharyngitis, pleurisy, sinusitis, influenza, respiratory
 syncytial virus infection, herpes infection, HIV infection, hepatitis B virus infection,
 hepatitis C virus infection, disseminated bacteremia, Dengue fever, candidiasis,
 15 malaria, filariasis, amebiasis, hydatid cysts, burns, dermatitis, dermatomyositis,
 sunburn, urticaria, warts, wheals, vasculitis, angiitis, endocarditis, arteritis,
 atherosclerosis, thrombophlebitis, pericarditis, myocarditis, myocardial ischemia,
 periarteritis nodosa, rheumatic fever, Alzheimer's disease, coeliac disease, congestive
 heart failure, adult respiratory distress syndrome, meningitis, encephalitis, multiple
 20 sclerosis, cerebral infarction, cerebral embolism, Guillame-Barre syndrome, neuritis,
 neuralgia, spinal cord injury, paralysis, uveitis, arthritides, arthralgias, osteomyelitis,
 fasciitis, Paget's disease, gout, periodontal disease, rheumatoid arthritis, synovitis,
 myasthenia gravis, thyroiditis, systemic lupus erythematosus, Goodpasture's
 syndrome, Behcets's syndrome, allograft rejection, graft-versus-host disease, Type I
 25 diabetes, ankylosing spondylitis, Berger's disease, Type I diabetes, ankylosing
 spondylitis, Retier's syndrome, or Hodgkins disease. In more preferred
 embodiments, the condition is appendicitis, peptic, gastric or duodenal ulcers,
 peritonitis, pancreatitis, ulcerative, pseudomembranous, acute or ischemic colitis,
 hepatitis, Crohn's disease, asthma, allergy, anaphylactic shock, organ ischemia,
 30 reperfusion injury, organ necrosis, hay fever, sepsis, septicemia, endotoxic shock,

cachexia, septic abortion, disseminated bacteremia, burns, Alzheimer's disease, coeliac disease, congestive heart failure, adult respiratory distress syndrome, cerebral infarction, cerebral embolism, spinal cord injury, paralysis, allograft rejection and graft-versus-host disease.

5

55. The method of Claim 54 wherein the disorder is selected from the group consisting of peritonitis, pancreatitis, ulcerative colitis, Crohn's disease, asthma, organ ischemia, reperfusion ischemia, sepsis, cachexia, burns, myocardial ischemia, adult respiratory distress syndrome, multiple sclerosis, rheumatoid arthritis, systemic lupus erythematosus, chronic obstructive pulmonary disease, psoriasis, Behcet's syndrome, allograft rejection and graft-versus-host disease.





10

56. The method of Claim 55 wherein the disorder is sepsis.

ANTI-INFLAMMATORY COMPOUNDS

ABSTRACT OF THE DISCLOSURE

- Compounds of Formula (I), pharmaceutical compositions comprising compounds of
5 Formulae (I a) or (VII) and a method of treating a subject with an inflammatory cytokine-mediated disorder comprising administering to the subject a compound of Formulae (I a) or (VIIa).



- 10 The variables of Formulae (I), (I a), (VII) and (VII a) are described herein.




FIG. 1




FIG. 2

Compound	IC ₅₀ (μM)
Nicotine	~ 1
VI f	< 1
VI b	~ 1
XII b	~ 1
VI e	~ 1
VI g	> 10
XII a	> 100

FIG. 3

	Carrier	Compound (VI f)
Day-1	13	13
Day 0	13	13
Day 1	9	12
Day 2	9	12
Day 3	9	12
Day 4	9	12
Day 5	7	12
Day 6	7	12
Day 7	7	12
Day 8	7	11
Day 9	7	11
Day 10	7	11
Day 11	7	11
Day 12	7	11
Day 13	7	11

FIG. 4




FIG. 5




FIG. 6

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/US04/032986

International filing date: 07 October 2004 (07.10.2004)

Document type: Certified copy of priority document

Document details: Country/Office: US
Number: 60/560,719
Filing date: 07 October 2003 (07.10.2003)

Date of receipt at the International Bureau: 18 November 2004 (18.11.2004)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse